

CMS Clinical Quality Measures: EHR Incentive Program

Core			
Measure	Developer	Title	Description
0013	AMA	Hypertension: Blood Pressure Measurement	Percentage of patient visits for patients aged 18 years and older with a diagnosis of hypertension who have been seen for at least 2 office visits, with blood pressure (BP) recorded.
0028a	AMA	Preventive Care and Screening Measure Pair: a.Tobacco Use Assessment	Percentage of patients aged 18 years or older who have been seen for at least 2 office visits, who were queried about tobacco use one or more times within 24 months.
0028b	AMA	Preventive Care and Screening Measure Pair: b.Tobacco Cessation Intervention	Percentage of patients aged 18 years and older identified as tobacco users within the past 24 months who received cessation intervention.
0421	QIP	Adult Weight Screening and Follow-Up	Percentage of patients aged 18 years and older with a calculated BMI in the past six months or during the current visit documented in the medical record AND if the most recent BMI is outside parameters, a follow-up plan is documented.
AlternateCore			
Measure	Developer	Title	Description
0024	NCQA	Weight Assessment and Counseling for Children and Adolescents	The percentage of patients 2-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of BMI percentile documentation, counseling for nutrition and counseling for physical activity during the measurement year
0041	AMA	Preventive Care and Screening: Influenza Immunization for Patients ≥ 50 Years Old	Percentage of patients aged 50 years and older who received an influenza immunization during the flu season (September through February).
0038	NCQA	Childhood immunization Status	The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); two H influenza type B (HiB); three hepatitis B (Hep B), one chicken pox (VZV); four pneumococcal conjugate (PCV); two hepatitis A (Hep A); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. The measure calculates a rate for each vaccine and two separate combination rates.